1等学療

100719. CINNAMOMUM BURMANNT | 100724 to 100798—Continued. Blume. Lauraceae.

rom Sumatra. Seeds presented by Edward Jacobson, Fort de Kock. Received July 9, 1932. From Sumatra.

An evergreen tree, native to Java, with shining green, ovate-oblong, papery leaves of variable size, usually 2 to 6 inches long, emitting a cinnamon

For previous introduction see 94258.

100720. LIGUSTRUM JAPONICUM ROTUN-Blume (L. coriaceum DIFOLIUM Carr.). Oleaceae. Dwarf privet.

From South Carolina. Cuttings presented by the Pinehurst Nursery, Summerville. Received Pinehurst 1 July 14, 1932.

A compact form of the Japanese privet, up to 6 feet high, with crowded, suborbicular, lustrous, evergreen leaves 2 to 3 inches long, and dense panicles, 2 to 4 inches long, of small white flowers followed by small, blue-black, subglobose fruits.

100721. Sesamum orientale L. Pedaliaceae. Sesame.

From Palestine. Seeds presented by the director of Agriculture, Forests, and Fisheries, Jerusalem. Received July 13, 1932.

From the Government Experiment Station at Beisan. One of the best types grown.

100722. CARYOPHYLLUS JAMBOS Myr-Stokes (Eugenia jambos L.). Rose-apple.

From the West Indies. Seeds presented by Stephen Hawels, Mount Joy, Dominica, British West Indies. Received June 19, 1931. Numbered in

A medium-sized handsome tree, native to India and the Malay Peninsula, but cultivated in many tropical countries for the edible, fragrant, pinkish fruits which are about the size of a hen's egg, of a sweetish acid taste, and said to be used as preserves. It thrives best in moist regions at elevations up to 3,000 feet, preferring a deep rich soil; it is propagated by seed.

For previous introduction see 93358.

100723. MEDICAGO SATIVA L. Faba-Alfalfa. ceae.

rom Turkey. Seeds presented by Charles E. Allen, American consul, Istanbul, through the State Department. Received July 20, 1932. From Turkey.

Turkish alfalfa, introduced for the use of Department specialists interested in forage crops.

100724 to 100798. Solanum Tubero-Solanaceae. Potato. SUM.

From South America. Tubers collected by H. G. MacMillan and C. O. Erlanson, Bureau of Plant Industry. Received July 12, 1932.

100724 and 100725. Collected in the region of Cuzco, Peru, May 25 and 26, 1932, at 11,000 feet altitude.

100724. No. 382. Medium-sized, elongated, gray-purple tubers, with heavy-browed, shallow eyes and white flesh with purple vascular ring.

100725. No. 383. Medium-sized, ovate, flat-tened tubers, with white skin mottled with red, shallow eyes and white flesh.

100726 to 100735. Collected June 5, 1932, at Cochabamba, Bolivia, at 8,300 feet altitude.

100726. No. 384. Round, purple and red mot-tled tubers, which have yellow-white flesh and deep eyes with pink buds.

100727. No. 385. Round white tubers tinted with pink, deep open eyes and white flesh.

100798. No. 386. Long, slim, curved or curled, twisted tubers, with thick dark skin, deep closed eyes, and white flesh with the center

100729. No. 387. Long, cylindrical tubers, with thick gray-black skin which becomes purple when cut, shallow eyes, and yellow-white

100730. No. 388. Long, cylindrical, pink tubers with deep white eyes.

0731. No. 389. Round, pink and white tubers, flat at the bud end, with open, medium deep eyes and white flesh. 100731. No.

100732. No. 390. Large, long, cylindrical, tan-white tubers, with deep eyes and lightly mottled white flesh.

100733. No. 391. Round white tubers, some spotted with pink, with open flat eyes, some showing pink, and white flesh.

100734. No. 392. Small, round yellow tubers spotted with red, shallow eyes with long eyebrows, and flesh yellow at vascular region, fading to white at center of potato.

100735. No. 393. Round, small to medium-sized, dark-purple tubers, with deep open eyes full of red buds, and yellow-white fiesh with purple spots in the vascular ring.

100736 to 100798. Collected from June 9 to June 19, near La Paz, Bolivia, at 12,500 feet altitude.

100736. No. 401. Chaclo Chaclo. small pink tubers generally pointed at the bud end, with numerous deep, straight eyes and yellow-white flesh flecked with red inside the vascular ring.

20737. No. 402. Cunduma. Small, round, black tubers, except around the numerous deep eyes, where they are yellowish white, and yellowish flesh with purple flecks, heaviest at the stem end. This is locally known as a sweetpotato, papa dulce.

100738. No. 403. Rinkel. Round, medium-sized tubers, yellowish to dirty white and black around the stem; the center of the medium to deep eyes is white, surrounded by black, and the flesh is clear yellow. Known locally as a sweetpotato, papa dulce.

00739. No. 404. Chapallo. Small-sized, round brown-tan tubers with clear yellowish flesh and deep, brown-tan eyes with bulging eye-100739. No. 404.

00740. No. 405. Thuila Nimilla. Mediumsized, round, pink tubers with deep, pink 100740. No. 405. eyes and white flesh.

100741. No. 406. Cholo Colorado. Small-sized round red tubers with eyes exceptionally deep, open at the stem end, deeply notched at the bud end. The yellowish flesh is flecked throughout with red.

100742. No. 407. Follhaco. Tubers medium-sized, round, eyes medium deep, bulging between the eyes, skin red, white below the eyes. Flesh clear white.

10743. No. 408. Choquefrit. Round, medium-sized, white tubers, tinted pink at the stem end, medium to shallow eyes with very long eyebrows and clear white flesh. This is known as a bitter variety. 100743. No. 408.

Pidriaja. 100744. No. 409. Medium-sized. round, flat. pink to purple tubers with white flesh. The deep open eyes have short eyebrows and vary in color as the tuber does.

100745. No. 410. Pulo Colorado. Long, cylindrical, medium-sized, red tubers with thick skin, which is slightly darker at the few shallow eyes; the flesh is yellowish white.